

VERIFICATION OF TRANSLATION

I undersigned, Mr. Norberto VESGA

Of C. Consell de Cent, 322; 08007 Barcelona; Spain


declare as follows:

1. That I am well acquainted with both the English and Spanish languages, and
2. That the attached document is a true and correct translation into English made by me to the best of my knowledge and belief of:

The Spanish Patent of Invention n° 200301340 filed on May 29, 2003

Barcelona, November 18, 2005

Signature of Translator: _____

A handwritten signature in black ink, appearing to read "Norberto Vesga", is written over a horizontal line. The signature is stylized with a large loop at the beginning and a long horizontal stroke at the end.

OFFICIAL CERTIFICATE

I hereby certify that the annexed documents are an exact copy of the PATENT OF INVENTION number 200301340, according to the grant issued on date May 29, 2003

Madrid, May 10, 2004

The Director of the Patents Department
and Technological Information

(signature)
CARMEN LENCE REIJA

[Seal of the
Spanish Patents and
Trademarks Office]

SPANISH PATENTS AND TRADEMARKS OFFICE

[Stamp bearing: Files-Association of Industrial Property Agents]

APPLICATION NUMBER
P200301340

DATE AND HOUR OF FILING AT THE O.E.P.M.
29 May 2003

INSTANCE OF APPLICATION FOR:

☒ PATENT OF INVENTION

☐ UTILITY MODEL

- (2)
- ☐ ADDITIONAL APPLICATION
- ☐ DIVISIONAL APPLICATION
- ☐ CHANGE OF MODALITY
- ☐ TRANSFORMATION
- ☐ EUROPEAN APPLICATION

(3) PRINCIPAL OR ORIGINAL FILE

MODALITY

APPLICATION NUMBER

FILING DATE

MODALITY

APPLICATION NUMBER

FILING DATE

DATE AND HOUR OF FILING IN A PLACE
OTHER THAN THE O.E.P.M.

(4) PLACE OF FILING CODE

BARCELONA 08

(5) APPLICANT (S) SURNAME OR JURIDICAL DENOMINATION	NAME	NATIONALITY	COUNTRY CODE	D.N.I.
QUÍMICA SINTETICA, S.A.		SPANISH	ES	A28008167

(5) PARTICULARS OF THE FIRST APPLICANT

ADDRESS: C. de l'Acer, 24

TOWN: BARCELONA

PROVINCE: BARCELONA

COUNTRY OF RESIDENCE: SPAIN

NATIONALITY: Spanish

PHONE

POST CODE 08004

COUNTRY CODE [ES]

NATION CODE [ES]

(7) INVENTOR (S)	(8) <input type="checkbox"/> THE APPLICANT IS THE INVENTOR	(9) WAY OF OBTENTION OF RIGHTS
	<input checked="" type="checkbox"/> THE APPLICANT IS NOT THE INVENTOR OR SOLE INVENTOR	<input checked="" type="checkbox"/> EMPLOYER (S) <input type="checkbox"/> CONTRACT <input type="checkbox"/> SUCCESSION

SURNAMES	NAME	NATIONALITY	CODE
COSME GOMEZ	ANTONIO	SPANISH	ES
PALOMO NICOLAU	FRANCISCO EUGENIO	SPANISH	ES

(9) TITLE OF THE INVENTION

ADDITION SALTS OF AZITHROMYCIN AND CITRIC ACID AND PROCESS FOR PREPARING THEM.

(11) MICROBIOLOGICAL DEPOSIT ☐ YES ☐ NO

(12) OFFICIAL EXHIBITIONS: PLACE PLACE DATE

(13) PRIORITY DECLARATIONS	
COUNTRY OF ORIGIN	CODE NUMBER DATE

(14) THE APPLICANTS REQUEST THE EXEMPTION OF THE PAYMENT OF TAXES PROVIDED IN ART. 162 L.P. ☐ YES ☐ NO

(15) REPRESENTATIVE	SURNAMES: PONTI SALES	NAME CODE
		Adelaida [388/3]
ADDRESS	TOWN	PROVINCE POSTAL
C. Consell de Cent, 322	Barcelona	CODE Barcelona [08007]

(16) LIST OF ANNEXED DOCUMENTS	SIGNATURE OF THE APPLICANT OR REPRESENTATIVE
<input checked="" type="checkbox"/> SPECIFICATION Nº OF PAGES 11 <input checked="" type="checkbox"/> CLAIMS Nº OF PAGES 27 <input checked="" type="checkbox"/> DRAWINGS Nº OF PAGES 15 <input type="checkbox"/> SEQUENCE LISTING 0 <input checked="" type="checkbox"/> ABSTRACT <input type="checkbox"/> PRIORITY DOCUMENT <input type="checkbox"/> TRANSLATION OF PRIORITY DOCUMENT	Adelaida Ponti Sales Associate no. 320
<input checked="" type="checkbox"/> REPRESENTATION FORM <input type="checkbox"/> PROOFS <input checked="" type="checkbox"/> RECEIPT OF PAYMENT OF TAXES <input type="checkbox"/> SHEET OF COMPLEMENTARY INFORMATION <input checked="" type="checkbox"/> OTHER Magnetic Support and Inventor's Declaration	SIGNATURE OF THE OFFICIAL

NOTIFICATION OF PAYMENT OF THE GRANT FEES

To the Director of the Spanish Patents and Trademarks Office

informacion@oepm.es

www.oepm.es

C/Panamá 1, 28071 MADRID

ABSTRACT AND GRAPHIC

ABSTRACT

ADDITION SALTS OF AZITHROMYCIN AND CITRIC ACID AND PROCESS FOR PREPARING THEM

Said addition salts have a molar ratio between azithromycin and citric acid such as to provide a pH comprised between 4.0 and 8.0 in a 10% aqueous solution.

The process for preparing said salts comprises: a) dissolving azithromycin in a solvent or mixture of solvents; b) adding citric acid-; and c) isolating the product obtained by crystallisation.

The addition salts of azithromycin and citric acid are stable and soluble in aqueous medium, being useful antibacterial and antiprotozoan agents

GRAPHIC

LAP9 Rec'd PCT/PTO 29 NOV 2005

FILING DATE : 29 MAY 2003

NUMBER DATE COUNTRY	
APPLICANT(S) QUÍMICA SINTÉTICA, S.A. ADDRESS: C. de l'Acer, 24 NATIONALITY: SPANISH 08004 BARCELONA, SPAIN	
INVENTOR(S) ANTONIO COSME GOMEZ, FRANCISCO EUGENIO PALOMO NICOLAU	
INT. CL TITLE OF THE INVENTION ADDITION SALTS OF AZITHROMYCIN AND CITRIC ACID AND PROCESS FOR PREPARING THEM.	GRAPHIC
ABSTRACT ADDITION SALTS OF AZITHROMYCIN AND CITRIC ACID AND PROCESS FOR PREPARING THEM. Said addition salts have a molar ratio between azithromycin and citric acid such as to provide a pH comprised between 4.0 and 8.0 in a 10% aqueous solution. The process for preparing said salts comprises: a) dissolving azithromycin in a solvent or mixture of solvents; b) adding citric acid-; and c) isolating the product obtained by crystallisation. The addition salts of azithromycin and citric acid are stable and soluble in aqueous medium, being useful antibacterial and antiprotozoan agents	


**Addition salts of azithromycin and citric acid and
process for preparing them**

Field of the invention

This invention relates to new addition salts of
5 azithromycin and citric acid, their preparation, their use
in pharmaceutical compositions and the aqueous or water-
alcohol solutions containing them, as well.

Background of the invention

Azithromycin or 9-deoxo-9a-aza-9a-methyl -9a-
10 homoerythromycin A:



is a broad-spectrum antibacterial agent which was
described and patented by Sour Pliva in Yugoslavian patent
application YU 000592 of 06/03/81, priority claimed in the
15 equivalent American patent US 4.517.359.

On the other hand, European patent EP 298650
describes azithromycin monohydrate and azithromycin
dihydrate. Chinese patents CN 1123279A, CN 1157824A and CN
20 1205338A, describe methods for preparing azithromycin
salts with organic and inorganic acids. The publication J.

Chem. Research (M), 1988,1239-1261; *J. Chem. Research (S)*, 1988,152-153 describe azithromycin dihydrochloride, dihydroiodide, diacetate, diaspartate, diglucoheptonate and dilactobionate. Patent application WO 00/32203
5 discloses azithromycin ethanolate and European patent application EP 984020 discloses an isopropanol caltrate of azithromycin monohydrate. Patent application WO 02/094843 discloses various crystalline forms of azithromycin, characterised by the carbon 13 nuclear magnetic resonance
10 spectrum (^{13}C -NMR) and the X-ray diffraction spectrum.

It is known that azithromycin is not stable in an aqueous acid medium, and furthermore base azithromycin is very insoluble in water.

There is therefore a need for providing new acid
15 addition salts of azithromycin that are soluble in aqueous medium while at the same time having suitable stability properties in solid phase and in solution.

Brief description of the invention

The object of this invention is to provide new
20 addition salts of azithromycin and citric acid soluble in aqueous medium while at the same time having suitable stability properties in solid phase and in solution.

A further object of this invention is to provide a process that is useful for preparing such salts and their
25 use for therapeutic purposes.

Brief description of the figures

Figure 1 shows the X-ray diffraction spectrum of azithromycin hydrogen citrate.

30 Figure 2 shows the carbon 13 nuclear magnetic resonance spectrum (^{13}C -NMR) of azithromycin hydrogen citrate in solid state.

Figure 3 shows the IR spectrum of azithromycin hydrogen citrate, recorded on KBr tablet.

Figure 4 shows the X-ray diffraction spectrum of azithromycin hydrogen citrate.

Figure 5 shows the carbon 13 nuclear magnetic resonance spectrum (^{13}C -NMR) of azithromycin hydrogen citrate in solid state.

Figure 6 shows the IR spectrum of azithromycin hydrogen citrate, recorded on KBr tablet.

Figure 7 shows the X-ray diffraction spectrum of azithromycin hydrogen citrate.

10 Figure 8 shows the carbon 13 nuclear magnetic resonance spectrum (^{13}C -NMR) of azithromycin hydrogen citrate in solid state.

Figure 9 shows the IR spectrum of azithromycin hydrogen citrate, recorded on KBr tablet.

15 Figure 10 shows the X-ray diffraction spectrum of azithromycin citrate.

Figure 11 shows the carbon 13 nuclear magnetic resonance spectrum (^{13}C -NMR) of azithromycin citrate in solid state.

20 Figure 12 shows the IR spectrum of azithromycin citrate, recorded on KBr tablet.

Figure 13 shows the X-ray diffraction spectrum of azithromycin citrate.

25 Figure 14 shows the carbon 13 nuclear magnetic resonance spectrum (^{13}C -NMR) of azithromycin citrate in solid state.

Figure 15 shows the IR spectrum of azithromycin citrate, recorded on KBr tablet.

30

Detailed description of the invention

Surprisingly, the authors of this invention have found new addition salts of azithromycin and citric acid which show good solubility in aqueous medium and good
35 stability properties in solid phase and in solution.

In a first aspect, this invention relates to a new addition salt of azithromycin and citric acid, the molar ratio between the azithromycin and the citric acid being such as to provide a pH between 4.0 and 8.0 in a 10% aqueous solution.

In one embodiment of the invention, said salt is azithromycin hydrogen citrate, which is characterised in that it has a molar ratio of azithromycin and citric acid such as to provide a pH between 4.0 and 6.0 in 10% aqueous solution.

For the purposes of the present invention and except where expressly stated otherwise, the percentage of the addition salt of azithromycin and citric acid in aqueous solution is expressed in weight/weight or weight/volume.

Preferably, the azithromycin hydrogen citrate salt contains up to 8% water, more preferably up to 6%, under relative humidity conditions of 40%.

More preferably still, said azithromycin hydrogen citrate further contains up to 3% of residual solvent.

Advantageously, said azithromycin hydrogen citrate is characterised in that it has a molar ratio of azithromycin and citric acid close to the stoichiometric ratio that provides a pH of 5 in a 10% aqueous solution.

In a second embodiment of the invention, said salt is azithromycin citrate, which is characterised by having a molar ratio of azithromycin and citric acid such as to provide a pH between 6.0 and 8.0 in 10% aqueous solution.

Preferably, the azithromycin citrate salt contains up to 8% water, and more preferably still up to 6%, under relative humidity conditions of 40%.

More preferably still, the azithromycin citrate further contains up to 3% of residual solvent.

Advantageously, said azithromycin citrate has a molar ratio of azithromycin and citric acid of 3:2.

Also advantageously, the azithromycin citrate, in accordance with one preferable embodiment of the present invention, is in amorphous form.

The azithromycin citrate in accordance with one
5 embodiment of the invention is characterised in that it has a chemical combination of one molecule of azithromycin per 2/3 of a molecule of citric acid (chemically, 3 moles of azithromycin and 2 moles of citric acid), resulting in a neutral salt in which the basic groups of azithromycin
10 (two equivalents) form a salt with the acid groups of the citric acid (3 equivalents).

The azithromycin citrate of the invention provides aqueous solutions up to 65% (w/w) at ambient temperature, with a pH between 6.8 and 7.5.

15 A second aspect of the invention is to provide a process for preparing an addition salt of azithromycin and citric acid, in accordance with the first aspect of this invention. Such process comprises: a) dissolving azithromycin in a solvent or mixture of solvents, b)
20 adding citric acid; and c) isolating the product obtained.

Citric acid or 2-Hidroxy-1,2,3-propanotricarboxylic acid is a carboxylic acid that has three COOH groups in its molecule.

Azithromycin has two nitrogen groups of basic nature
25 in its molecule and for the process of the invention can be used either in monohydrate or dihydrate form of azithromycin.

In one embodiment of the process of the invention, step (a) is carried out by dissolving azithromycin in
30 monohydrated form.

In another embodiment, step (a) is carried out by dissolving azithromycin in dihydrated form.

For the purposes of this invention, unless expressly stated otherwise, dissolving azithromycin in a solvent or
35 mixture of solvents should be understood to mean any

degree of dissolution, with total dissolution of the product at the start of the process being unnecessary.

The addition salt of azithromycin and citric acid can be prepared in practically any kind of solvent, although it is more difficult its preparation in solvents in which both molecules are insoluble (for example, in toluene or heptane). The following can be used as solvents: water; linear or branched C₁-C₆ aliphatic alcohols, such as methanol, ethanol, n-propanol, isopropanol, n-butanol, etc.; cyclic aliphatic alcohols, such as cyclohexanol; diols, such as ethylene glycol, 1,2-propylene glycol, 1,3-propanediol, 1,4-butanediol, etc.; linear or branched C₁-C₆ aliphatic ketones, such as acetone, methyl ethyl ketone, methyl isobutyl ketone, etc.; cyclic aliphatic ketones, such as cyclohexanone; short-chain aliphatic esters, such as methyl or ethyl acetate; short-chain aliphatic ethers, such as ethylic ether, isopropylic ether, etc.; cyclic aliphatic ethers, such as tetrahydrofuran and dioxane, or mixtures thereof.

In one embodiment of the process of the invention, the azithromycin hydrogen citrate salt is prepared by isolating the salt by means of crystallisation in step (c).

The following aspects, independently or together, are preferred in the preceding embodiment: the azithromycin is selected from the azithromycin monohydrate or dihydrate; the molar proportions of azithromycin and citric acid are close to the stoichiometric; the solvents are selected from alcohols, ketones, esters or ethers or mixtures thereof, preferably ethanol, acetone, methyl acetate or tetrahydrofuran or mixtures thereof; the crystallisation temperature is between 25°C and the solvent's reflux temperature; and the mixture is cooled to a temperature between 0° C and 25°C before separating the crystals.

The X-ray diffraction, carbon 13 nuclear magnetic resonance (^{13}C -NMR) in solid state and IR spectra serve to identify the azithromycin hydrogen citrate in accordance with the first aspect of the invention. See Figures 1 to 5 9.

In another embodiment of the process of the invention, the azithromycin citrate is prepared by adding an amount of citric acid in step (b) such that the molar ratio between the azithromycin and the citric acid is 3:2.

10 Advantageously, when the azithromycin citrate is prepared, the salt is isolated in step (c) by eliminating the solvent.

The following aspects, independently or together, are preferred in the preceding embodiment: the azithromycin is 15 selected from the azithromycin monohydrate or dihydrate; the solvents are selected from water, alcohols, ketones, esters or ethers, or mixtures thereof, preferably water, ethanol, acetone, methyl acetate or tetrahydrofuran, or mixtures thereof.

20 The X-ray diffraction, carbon 13 nuclear magnetic resonance (^{13}C -NMR) in solid state and IR spectra serve to identify the azithromycin citrate produced in accordance with the invention. See Figures 10 to 15.

The new aqueous-medium-soluble addition salts of 25 azithromycin and citric acid of the invention that have suitable stability characteristics in solid phase and in solution are useful as antibacterial and antiprotozoans. They can be administered orally, parenterally, topically or rectally in the treatment or prevention of infections 30 caused by bacteria or protozoa.

The new addition salts of azithromycin and citric acid of the invention are particularly useful in the preparation of aqueous or water-alcohol solutions of azithromycin containing up to 65% of the salt, providing a 35 pH between 4 and 8, stable and not suffering from chemical

degradation of azithromycin.

For a better understanding of all that has been described there follow some examples which show, 5 schematically and solely by way of non-restrictive example, some embodiments of the invention.

Examples

10 Example 1. Preparation of azithromycin hydrogen citrate

20g of azithromycin are added to 100ml of acetone (water content according to Karl-Fisher of 1 to 5%), the mixture is stirred at ambient temperature until dissolved. 5.35g of citric acid are added and the mixture is heated 15 at reflux. It is then cooled to 0-5°C, filtered, washed with acetone and dried under vacuum at 40°C to yield 22.4g of azithromycin hydrogen citrate (water content according to Karl-Fisher of 1.2% and acetone content less than 0.5%). The azithromycin content determined by HPLC is 80% 20 and the citric acid content by electrometric titration is 20%, corresponding to the stoichiometric ratio of the azithromycin hydrogen citrate. The salt can contain up to 8% water depending on the drying method (by vacuum, fluidised bed, static), but is preferably 6%, under 25 relative humidity conditions of 40%. Figures 1, 2 and 3 show the X-ray diffraction spectrum, the carbon 13 nuclear magnetic resonance spectrum (¹³C-NMR) in solid state and the IR spectrum, recorded on KBr tablet, respectively.

30 Example 2. Preparation of azithromycin hydrogen citrate

20g of azithromycin dihydrate and 3.5g of citric acid monohydrate are added to 50 ml of methyl acetate. This is heated at reflux, cooled to ambient temperature, filtered, washed with methyl acetate and dried under vacuum at 40°C. 35 Figures 4, 5, and 6 show the X-ray diffraction spectrum,

the carbon 13 nuclear magnetic resonance spectrum (^{13}C -NMR) in solid state and the IR spectrum, recorded on KBr tablet, respectively.

5 Example 3. Preparation of azithromycin hydrogen citrate

Following the procedure set out in example 2 and replacing the methyl acetate by tetrahydrofuran, azithromycin hydrogen citrate is obtained. Figures 7, 8 and 9 show the X-ray diffraction spectrum, the carbon 13
10 nuclear magnetic resonance spectrum (^{13}C -NMR) in solid state and the IR spectrum, recorded on KBr tablet, respectively.

Example 4. Preparation of azithromycin citrate

15 20g of azithromycin dihydrate and 3.5g of citric acid monohydrate are dissolved at ambient temperature in 50 ml of ethanol, filtered and the solvent is evaporated at low pressure. 24.9g of a white solid is obtained, containing up to 2.0% of ethanol and up to 7% of water. The X-ray
20 diffraction spectrum confirms that it is an amorphous product (Fig. 10). Figures 10, 11 and 12 show the X-ray diffraction spectrum, the carbon 13 nuclear magnetic resonance spectrum (^{13}C -NMR) in solid state and the IR spectrum, recorded on KBr tablet, respectively.

25

Example 5. Preparation of azithromycin citrate

20g of azithromycin dihydrate and 3.5g of citric acid are added to 50 ml of water. The mixture is stirred at ambient temperature and the insoluble material
30 filtered. The solution is concentrated at low pressure to a KF of around 5%, yielding 23.1g of azithromycin citrate. Figures XIII, XIV and XV show the X-ray diffraction spectrum, the carbon 13 nuclear magnetic resonance spectrum (^{13}C -NMR) in solid state and the IR spectrum,
35 recorded on KBr tablet, respectively.

Example 6. Preparation of azithromycin citrate solutions

Azithromycin citrate solutions are prepared by adding 20g of azithromycin, 3.5g of citric acid and the
5 corresponding amount of water (35 to 94g of water), stirring at ambient temperature for a time ranging between 30 and 60 minutes, and finally filtering to remove insoluble material. The solution is stable at ambient temperature.

10 Although specific embodiments of this invention have been described and shown, it is obvious that one skilled in the art could introduce variants and alterations, or replace details by others that are technically equivalent without departing from the
15 protection defined by the attached claims.

|

CLAIMS

1. Addition salt of azithromycin and citric acid,
5 in which the molar ratio between the azithromycin and the
citric acid is such as to provide a pH between 4.0 and
8.0 in a 10% aqueous solution.

2. Addition salt of azithromycin according to Claim
10 1, characterised in that it is azithromycin hydrogen
citrate.

3. Addition salt of azithromycin according to Claim
1, characterised in that it is azithromycin citrate.
15

4. Addition salt of azithromycin according to Claim
1, characterised in that it includes up to 8% water.

5. Addition salt of azithromycin according to Claim 4,
20 characterised in that it further includes up to 6% by
weight of water.

6. Addition salt of azithromycin according to Claim
1, which further contains up to 3% of residual matter.
25

7. Addition salt of azithromycin according to Claims
1 and 2, characterised in that the salt has a molar ratio
of azithromycin and citric acid such that it provides a pH
between 4.0 and 6.0 in a 10% aqueous solution.
30

8. Addition salt of azithromycin according to Claims
1 and 3, characterised in that the salt has a molar ratio
of azithromycin and citric acid such as to provide a pH
between 6.0 and 8.0 in a 10% aqueous solution.
35

9. Addition salt of azithromycin according to Claim 2 and 4, characterised in that with the molar ratio of azithromycin and citric acid being 1:1 a pH of 5 is provided in a 10% aqueous solution.

5

10. Addition salt of azithromycin according to Claims 3 and 5, characterised in that the molar ratio of azithromycin and citric acid is 3:2.

10 11. Addition salt of azithromycin according to Claim 3, characterised in that it is in an amorphous state.

12. Process for preparing an addition salt of azithromycin and citric acid according to Claim 1,
15 characterised in that it comprises:

- a) dissolving azithromycin in a solvent or mixture of solvents;
- b) adding citric acid; and
- c) isolating the product obtained.

20

13. Process according to Claim 12, characterised in that azithromycin is dissolved in monohydrated form in step (a).

25 14. Process according to Claim 12, characterised in that azithromycin is dissolved in dihydrated form in step (a).

15. Process according to Claim 12 characterised in
30 that the solvent is selected from: water; the linear or branched C₁-C₆ aliphatic alcohols, such as methanol, ethanol, n-propanol, isopropanol, n-butanol; cyclic aliphatic alcohols, such as cyclohexanol; diols, such as ethylene glycol, 1,2-propylene glycol, 1,3-propanediol,
35 1,4-butanediol; linear or branched C₁-C₆ aliphatic

ketones, such as acetone, methyl ethyl ketone, methyl isobutyl ketone; cyclic aliphatic ketones, such as cyclohexanone; short-chain aliphatic esters, such as ethyl acetate; short-chain aliphatic ethers, such as ethyl ether, isopropyl ether, etc.; cyclic aliphatic ethers, such as tetrahydrofuran and dioxane, or mixtures thereof.

16. Process according to Claim 15, characterised in that the azithromycin monohydrate or dihydrate is dissolved; the solvent is selected from water, alcohols, ketones, esters or ethers, or mixtures thereof, preferably water, ethanol, acetone, methyl acetate or tetrahydrofuran, or mixtures thereof.

17. Process according to any of Claims 12 to 16, for the preparation of azithromycin hydrogen citrate, characterised in that an amount of citric acid is added in step (b) such that the molar ratio between the azithromycin and the citric acid is close to the stoichiometric.

18. Process according to any of Claims 12 to 17, for the preparation of azithromycin hydrogen citrate, characterised in that in step (c) the salt is isolated by crystallisation.

19. Process according to Claim 18, characterised in that step c) comprises:

c-i) crystallising at a crystallisation temperature between 25°C and the solvent's reflux temperature; and
c-ii) cooling the mixture at a temperature between 0°C and 25°C, before separating the crystals.

20. Process according to any of Claims 12 to 17, for the preparation of azithromycin citrate, characterised in

that an amount of citric acid is added in step b) such that the molar ratio between the azithromycin and the citric acid is 3:2.

5 21. Process according to any of Claims 12 to 17 and 20, characterised in that for the preparation of azithromycin citrate, the salt is isolated by removing the solvent in step c).

10 22. Process for preparing solutions of an addition salt of azithromycin and citric acid according to Claim 1, in water or water-alcohol mixtures of up to 65%, which consists on: dissolving the azithromycin citrate in water and filtering the solution obtained.

15

23. Process for preparing solutions of at least one addition salt of azithromycin and citric acid, according to Claim 1, in water or water-alcohol mixtures of up to 65%, which consists on:

20 a) dissolving both components, azithromycin and citric acid, by stirring at ambient temperature; and
 b) filtering the solution obtained.

25 24. Azithromycin salt according to any of Claims 1 to 11 for use as an antibacterial agent.

25. Azithromycin salt according to any of Claims 1 to 11 for use as an antiprotozoan agent.

30

26. Use of an azithromycin salt according to any of Claims 1 to 11 for the manufacture of a medicament for the treatment of an infection caused by bacteria or protozoans.

35

27. Use of an azithromycin salt according to any of Claims 1 to 11 for the manufacture of a medicament for the prevention of an infection caused by bacteria or protozoans.

ABSTRACT

Said addition salts have a molar ratio between azithromycin and citric acid such as to provide a pH comprised between 4.0 and 8.0 in a 10% aqueous solution.

5 The process for preparing said salts comprises: a) dissolving azithromycin in a solvent or mixture of solvents; b) adding citric acid; and c) isolating the product obtained by crystallisation.

10 The addition salts of azithromycin and citric acid are stable and soluble in aqueous medium, being useful antibacterial and antiprotozoan agents.